AD	1		

Award Number: W81XWH-06-1-0693

TITLE: The Role of Smad4 on Tumorigenesis, ERK Pathway, and SERM Resistance in Estrogen Receptor Alpha-Positive Breast Carcinoma Cells

PRINCIPAL INVESTIGATOR: Robert Nichols

CONTRACTING ORGANIZATION: The University of Texas Health

Sciences Center

San Antonio, TX 78229-3900

REPORT DATE: September 2007

TYPE OF REPORT: ANNUAL SUMMARY

PREPARED FOR: U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for Public Release; Distribution Unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

a. REPORT	b. ABSTRACT	c. THIS PAGE		8	19b. TELEPHONE NUMBER (include area code)
16. SECURITY CLASS			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON USAMRMC
15. SUBJECT TERMS Predoctoral trai	ning, Breast cand	er research			
45 CHD IECT TERMS					
	·				
progress was sign award and was ev	ificantly hindered a entually deployed i	fter the PI was order	red to join National (The mentor, Dr. Luz	Guard for comb	ablished cell lines. However, the pat training two months after the asked by the DOD Program to
signaling pathways selective estrogen	s of TGF-beta/Sma receptor modulato	d4, ERK, and estrog rs. During the past y	gen in the regulation rear, the PI establish	of breast carcined various bre	nine the interaction among inogenesis and resistance to east cancer cell lines with
13. SUPPLEMENTAR	Y NOTES				
	ic Release; Distribu				
12 DISTRIBUTION / A	VAILABILITY STATEN	IFNT		N	NUMBER(S)
Fort Detrick, Mary	Research and Ma and 21702-5012	teriei Command		11. \$	SPONSOR/MONITOR'S REPORT
		IAME(S) AND ADDRESS	S(ES)	10. 5	SPONSOR/MONITOR'S ACRONYM(S)
The University San Antonio, TX		Sciences Cente	r		
	SANIZATION NAME(S)	, ,			ERFORMING ORGANIZATION REPORT UMBER
E-Mail: sunl@uths	csa.edu			5f. W	VORK UNIT NUMBER
Robert Nichols				5e. T	TASK NUMBER
6. AUTHOR(S)				5d. F	PROJECT NUMBER
					1XWH-06-1-0693 PROGRAM ELEMENT NUMBER
	J	s, ERK Pathway, an east Carcinoma Cell	d SERM Resistance in	5b. 0	GRANT NUMBER
4. TITLE AND SUBTIT	LE			5a. C	CONTRACT NUMBER
1. REPORT DATE (DE 29-SEP-2007	,	2. REPORT TYPE ANNUAL SUMMAR`	v		ATES COVERED (From - To) AUG 2006 - 29 AUG 2007
data needed, and completing a this burden to Department of D 4302. Respondents should be	and reviewing this collection of in Defense, Washington Headquart Daware that notwithstanding any	nformation. Send comments regarders Services, Directorate for Information	arding this burden estimate or any mation Operations and Reports in a shall be subject to any penalty f	y other aspect of this coll (0704-0188), 1215 Jeffer	ing existing data sources, gathering and maintaining the ection of information, including suggestions for reducing son Davis Highway, Suite 1204, Arlington, VA 22202- a collection of information if it does not display a currently

REPORT DOCUMENTATION PAGE

Form Approved OMB No. 0704-0188

Table of Contents

	<u>Page</u>
Introduction	4
Body	5
Key Research Accomplishments	6
Reportable Outcomes	7
Conclusion	8
References	
Appendices	

Introduction

It has been reported that 23% of breast tumors were smad4 negative, and 41% expressed low levels of smad4. Smad4 negative tumors had a longer disease free survival and better clinical outcome. Overexpression of Smad4 in estrogen receptor alpha positive breast carcinoma cells induces apoptosis via bim and bax intrinsic pathway. However, in estrogen receptor alpha negative breast carcinoma cells, Smad4 promotes epithelial-to-mesenchymal transdifferentiation and metastasis. For this predoctoral training application, we proposed to test the hypothesis that TGF-beta signaling mediated by smad4 positively regulates the ERK pathway, and negatively regulates estrogen receptor alpha, which inhibits the progression of tumorigenesis but promotes resistance to selective estrogen receptor modulators in MCF-7 and ZR-75-1 estrogen receptor alpha breast carcinoma cells.

Body

There were two tasks proposed for the first year of the award. In Task 1a, we will perform various assays, like MTT growth assay, Cell Cycle Analysis, Soft Agarose assay, migration and invasion assays, Apoptosis ELISA and Western Blot Analysis to determine the functional role of Smad4 on tumorigenesis. In Task 1b, we will orthotopically inject 5-6 week old female athymic nude mice and determine tumor burden and incidence, and euthanize mice at 10 weeks to determine the histological classification, distal metastasis, apoptosis, and perform immunohistochemistry. Task 1a was partially accomplished and Task 1b has not been started. The PI was successful in stably overexpressing a tetracycline-regulatable Smad4 in ZR-75-1 breast cancer cells that express little Smad4 and in stably knocking down the expression of Smad4 in MCF-7 breast cancer cells with a RNA interference approach. Knockdown of Smad4 in MCF-7 breast carcinoma cells increased the protein and mRNA levels of estrogen receptor alpha (ERa). Sensitivity to estrogen and raloxifene was also enhanced with the knockdown of Smad4. Overexpression of Smad4 in ZR-75-1 cells decreased the protein and mRNA levels of ERalpha. Sensitivity to estrogen and raloxifene was also decreased when Smad4 was overexpressed. Smad4 also induced ERK1/2 phosphorylation and apoptosis and senescence, which could be inhibited when treated with an ERK inhibitor, PD98059. The research was halted after the PI was ordered to join National Guard for combat training two months after the award and was eventually deployed in Iraq in May 2007.

KEY RESEARCH ACCOMPLISHMENTS

- 1. Generated Smad4-overexpressing stable ZR-75-1 cells.
- 2. Generated MCF-7 cells with reduced Smad4 expression by a Smad4 shRNA.
- 3. Demonstrated the regulation of ERa expression by Smad4 pathway.

REPORTABLE OUTCOMES
Cell lines: Smad4-overexpressing ZR-75-1 cells and Smad4-underexpressing MCF-7 cells

CONCLUSIONS

Smad4 is a major intracellular mediator of TGF-beta signaling. Our study indicates that Smad4 negatively regulates ERa. This is consistent with the observations that many ERa positive breast cancer cells are resistant to TGF-beta suggesting that loss or attenuation of TGF-beta signaling may be a major mechanism for the ERa-mediated breast carcinogenesis. These breast tumors are likely to express higher levels of ERa and more responsive to anti-estrogen therapy. On the other hand, some ERa positive breast cancer that retains TGF-beta sensitive may express a relatively low level of ERa and thus be more resistant to anti-estrogen therapy resulting in a poorer outcome. Therefore, our study may yield novel insights that will explain Smad4-assoicated poor clinical outcome and decreased disease-free survival of breast cancer patients.